

Bio Sulfur



REF. AG181004

World's first and largest production

Environmentally friendly

BioSulfur

ECOBIO HOLDINGS CO.,LTD.

World No.1 PREMIUM

Agriculture, Livestock, Aquaculture



Company Overview



*To transform the existing
To create the nonexisting
And forge sustainability!*



- CEO** Hyo-Soon Song
- Academic** Keimyung University, Environmentology
- 1989** Founding of EcoBio Holdings Co., Ltd.
- 2005** Chairman of Bio-division of New & Renewable Energy Association
- 2013** Received Presidential Prize for Renewable Energy Award
- 2015** Present Auditor of New Renewable Energy Association
- 2016** Current EcoBio Holdings CEO
- 2018** Korean Society of Clean Technology Vice-Chairman

Company Name: EcoBio Holdings Co., Ltd. (KOSDAQ 038870)

Corporate Registration Number : 122-88-01030

Headquarters : 5, Seoun-ro 26-gil, Seocho-gu, Seoul

Factory : 61, Geowo-ro, Seo-gu, Incheon

Tel. : 02)3483-2900 **Fax :** 02)3483-2929

Email : biosulfa@ecobio.co.kr

Homepage : www.ecobio.co.kr

- 2016** EcoBio Holdings Co., Ltd.
- 2013** Presidential Award for Excellence
- 2012** Awarded the highest prize by a company specializing in the recycling of waste energy resources - Minister of Environment
- 2008** Goldman Sachs U.K attracts \$ 28 million in foreign capital
- 2007** KOSDAQ Listed
- 2002** Joined Korea Renewable Energy Association
- 1989** Founding of EcoBio Holdings (previous TotalENS)

Benefits of Sulfur

Essential Nutrients

Important constituent of our bones, skin, and hair

Skincare

Plays an important role in skin cells and tissues

Constipation

Can help to improve constipation through diuretic action

Constitution

Improves muscles and bones
Prevents balding and promotes hair growth

Insulin and Hormones


Activates insulin hormones
Effective in controlling diabetes

Detoxification

Detoxifies carcinogenic and disease-causing sources

Effects of Sulfur on People

20th century – the age of
“Vitamins”



21st century – the age of
“BioSulfa”

On average, the human body has 0.2% sulfur in its weight and should be maintained at all times.
Taking 0.5 to 1mg of sulfur per kilogram of body weight is the key to good health.

Sulfur deficiency

Sulfur deficiency is a major cause of baldness, keratinization of nails, toenails, as well as skin aging
Due to sulfur's detoxification ability, it is also called the antidote of the 21st century.

Beauty Mineral

Sulfur is a main component of amino acids that make up enzymes that affect collagen, a molecule that maintains the elasticity of the skin.
It is effective enough to be called beauty minerals

Sulfur Effects on the human body

Detoxification

Detoxification ability of heavy metals, chemicals, various pesticides, yellow-dust, etc.
Helps in releasing heavy metals such as yellow-dust, car fumes, seasonings, pesticides etc. from the body

Other qualities

Suppresses increase in blood cholesterol
Improves decomposition of platelets and improves sperm activity
Has anti-cancer and inhibitory effect
Anti-parasitic activity

Effects of Sulfur animal feeds on livestock(1/2)

Meat Quality Improvement

Tested on meat quality of cow , pigs, ducks, and chickens.
High quality meat produced and approved
(Increase in sirloin cut size with 20% decrease in fat layer)

Collagen Proteins

Richer meat broth and improved meat texture through increase in collagen proteins

Cholesterol

Decrease in overall cholesterol and increase in unsaturated fatty acids

Endocrine-Disrupting hormones

Helps alleviate build-up of endocrine-disrupting hormones
Decreased build-up of heavy metals in muscles

Livestock Disease Control

Decreased livestock death due to increase in immunity, preventing disease outbreaks
(rate of gain increased by 5~67% in all livestock, increased effectiveness of feed by 3~5%)

Reduces need of antibiotic administration, excellent deodorizer

Effects of Sulfur animal feeds on livestock(2/2)

Odor Removal

When cooking, the fishy smell of meat is replaced by a savory scent

Oil solidification

Oil after cooking does not solidify and the taste of the meat does not change after cooling.

Meat Tenderness

The meat is becomes tender and dryness of breast meat is reduced.



Meat Quality

Animals that consume sulfur create a white broth when boiled for stock, creating a light and clean quality

Effective detoxification effect

Removes heavy metals, agrochemicals, and growth hormones

Advantages of BioSulfa to livestock



Nutrient	Carbohydrate	Protein	Mineral (Metal component)
Absorption type	Smallest size Monosaccharide	Amino acid	Smallest size Single molecule

The effect of sulfur as feed additive is closely related to the size of the sulfur incorporated.
A simple structure of sulfur compounds can be easily absorb to both animals and plants.

BioSulfa is highly probable for absorption by livestock,


Because of its small size (1 to 4 μ m) compared to conventional sulfur (chemical sulfur, 400 μ m)

Effect of Sulfur on Crops

- ✓ Improves crop growth through sterilization and disinfection action
- ✓ Improves nourishment supply through soil sterilization
- ✓ Improves resistant to disease and pests for environmentally friendly farming

Environmentally friendly farming = Increases income of farm, balanced development of industry

➔ **Healthy public**



Vision

Although sulfur has been used in agriculture since centuries ago, the toxicity of sulfur has prevented it from extensive use. However, Biosulfa produced by microbial metabolism is expected to attract worldwide attention as eco-friendly new agriculture. Biosulfa, which helps soil sterilization and crop growth as well as being highly efficient, is expected to accelerate the expansion of the eco-friendly agriculture.


What is BioSulfa?

Biologically produced!

BioSulfa

- ✓ Solves the toxicity problem of chemically produced sulfur.
- ✓ Biologically produced by microbial digestion (eco-friendly)
- ✓ Combines the benefits of sulfur and microorganisms
- ✓ Can use in agriculture, livestock, and aquaculture

※ Biosulfur production process within the Bioreactor



Thiobacillus breakdown of hydrogen sulfide by bacterial metabolism

Applications of BioSulfa

Agriculture

Soil improvement, winter control, foliar spray



Livestock

Development of animal feed



Aquaculture

Good replacement of currently used antibiotics



And much more!

Medicine, cosmetics, deodorizer, etc

BioSulfa is an ECO-Friendly Organic Product that is produced naturally by microorganisms and can be classified as an **Organic Sulfur Compound**.



- ☑ Production by microbial metabolism
- ☑ The presence of sulfur (S) and carbon (C) in the same molecule
- ☑ Eco-friendly product

※ The classification of organic sulfur compounds or inorganic sulfur compounds depends on coexistence of sulfur (S) and carbon (C) within the same molecule.

* Organic sulfur compounds : **C**+H+O+N+S / Inorganic sulfur compounds : **S**+O

Comparison between BioSulfur and chemical sulfur

Certified Organic Material
(Post No. 1-6-014)
(Certification No. 1-6-002)



VS



Category	Bio Sulfur (<i>suspended concentrate</i>)	Petrochemical Sulfur (solid)
Definition	Bio sulfur produced from microbial metabolism	Chemical sulfur derived from chemical reactions
pH	Slightly Alkaline (pH 8.5) suspension	Not soluble in water
Sulfur Conc.	40% ± 3% (suspension)	100% solid powder
Density	1.35 g/cm ³	1.95 ~ 2.26 g/cm ³
Particle Size	1~10 µm particles (high fungicidal effect)	400~600µm (low fungicidal effect)
Hydrophilicity	Pharmacological cocktail effect (Hydrophilic-like, can be mixed with other substances)	Not compatible with other chemicals (Highly hydrophobic, can not be mixed with other substances))
Characteristics	Naturally suspended state, easily suspended in water	Not soluble in water, needs caustic soda and surfactant for suspension
	User in organic fertilizers, pesticides, cosmetics and pharmaceuticals Harmless to insects	Used in chemical fertilizer Can cause metal corrosion, generates toxic gas and can reduce the lifespan of plastic materials

Market Value of BioSulfa





Of the entire **Sulfur** production in the world

Only **0.0001%** of it is **BioSulfa!**


High in scarcity value

- World's largest biosulfa production facility
- First domestic biosulfa production and commercialization
- The only domestic technology for production biosulfa






Agricultural BioSulfa (fungicide) → raw material for daily supplies → raw material for cosmetics → raw material for medicine



Used for Various Purposes



BioSulfa combines the benefits of microbial digestion and sulfur to create an eco-friendly product of the highest quality. Over 30,000 farmers have experienced its advantages

Global Market for Environment Friendly Fertilizer/Fodder

Zero Chemicals
Organic



Environment friendly

ZERO Pesticide

Preserve land and secure safe food by implementing pesticide-free

Production expansion

Specialized farming techniques

Competitive fertilizer and livestock industry

Strengthen competitiveness

Stable returning to farming

Contribute to economic revitalization of agricultural and fishery by stable cultivation

Problems of using existing chemical fertilizer

- ☑ Between 1960 and 2000, the use of chemical fertilizers increased about 800% globally.
- ☑ The use of chemical fertilizers is continuously increasing due to the success of chemical fertilizers such as nitrogen and phosphoric acid.
- ☑ Excessive acidification of soil due to the effect of acid rain by overspray and industrialization.
- ☑ The acidification of soil brought acidification of the plants, so that humans body is also affected
Most pathogens breed well in acid, so that pests outbreak on acidic soil and .
- ☑ Excessive nitrogen in chemical fertilizers generates large amounts of greenhouse gases, accelerating climate change. ("About 1/4 of total nitrous oxide emission, one of global warming material, is known as "results of agricultural activities using chemical fertilizers.")

The crisis of existing agriculture

Acidification of soil

- Soil becomes strongly acidic (pH 5.0 ~ 5.5) by excessive use of chemical fertilizer, pesticides, heavy metals in the dust.
- Organic and nutrient content is low. (Soil organic content : 2.4% in Korea)
- Zero chemical agriculture, organic farming, were tried, however productivity and wide availability are low. → It is not a fundamental way to solve the problem.

The aging of the agricultural population


- Aging of agriculture population is serious due to concentration of population and low birth rate.
- Agricultural population decreased continuously.

FTA

- The crisis of agriculture industry due to the liberalization of export and import of various agricultural products.
- Need to change into organic and qualified product.

Agricultural and chemical fertilizer usage (Korea)

< Usage of pesticides and chemicals >



- ▶ Serious soil acidification due to excessive fertilizer/pesticide usage.
- ▶ Need to find a way to solve soil acidification problem.

* Source: Ministry of Agriculture, Forestry and Livestock Foods

* Note: Chemical fertilizer usage is based on agricultural standards (excluding industrial use and export use)

* Usage of fertilizer and pesticide is classified by nationwide only.

Effect & Necessary of BioSulfa

Consumer

- Safe food
- Delicious food
- Fresh food
- Residual pesticide-free food
- Domestic agricultural products

Farmers

- Increase product quantity
- Continuous cultivation
- Increase income
- Production of high quality agricultural products
- Consumer satisfaction

Distributor

- Securing qualified agricultural products
- Extension of expiration date
- Safe product supply
- Increase profitability

Government


- Development of competitive agricultural technology
- Protection of Primary Industry from FTA
- Environmental protection
- Export of agricultural technology



How to use BioSulfa



+



=

On growing crops
Ratio (500-fold dilution)
Foliar treatment

- * Material for organic products
- * Hydrophilicity ensures safe use and mixing convenience

- * Provide essential ingredients for plants growth
- * Small particle size allows spray

※ Shake it sufficiently before use. Adjust spraying interval and concentration depending on soil, crop conditions and environment situation.



BioSulfa


Effects, Applications, and certification

Biosulfur has different properties to chemical sulfur,
making biosulfur unique to widely available sulfur.


Fungicidal effects of BioSulfa

Why So Effective?

- 1) Optimal Particle Size : 1~10 μm
- 2) Particle Size Optimal for Sterilizing Mold Spores
- 3) Germs and viruses are smaller than BioSulfa particles



1~10 μm







BioSulfa's fungicidal effect

- 1) Fungi reproduce through spores → can maintain spores through unfavorable conditions
- 2) Regular sulfur has larger particle size than fungi → can't prevent reproduction

Existing pesticides → Increase pesticide resistance

BioSulfa → highly effective → no build-up of resistance



PEOPLE



- Detoxify heavy metals
- Helps maintain skin elasticity
- Helps lower cholesterol and degrade platelets.
- Helps bones grow stronger
- Removes inflammation and kills germs
- Used to help treat diabetes
- Helps to alleviate constipation

LIVESTOCK



- Meat becomes tender
- Removal of unique odor
- High content of unsaturated fatty acids
- Slow oil solidification for fat
- Nutritious animal feed
- Good for digestion (helps stamina)

CROPS




- Improves crop growth through sterilization and disinfection action
- Improves nourishment supply through soil sterilization
- Improves resistant to disease and pests for environmentally friendly farming
- Contributes significantly in increasing farm income
- Winter control, soil sterilization, foliar spray etc.
- Improves color, scent, and taste of fruits

Safe to use in agriculture, livestock, and aquaculture

Sulfur pig, sulfur cows, sulfur ducks, sulfur chickens, sulfur eggs, sulfur deer, sulfur larvae and sulfur mudfish, etc.

The possibility of non-antibiotics in agriculture, livestock, and aquaculture begins with BioSulfa!



Comparison of Safety of BioSulfa and processed Sulfur



(units : ppm)

Category	Permissible Standard	BioSulfa <small>Taken from Korea Testing & Research Institute 2016.10.10 results</small>	Processed Sulfur <small>Taken from Konkuk University, College of Animal Bioscience & Technology</small>
S(Sulfur)	-	50.8%	98.34%
Pb(Lead)	3.0	Undetected	1.32
Cd(Cadmium)	3.0	Undetected	0.06
As(Arsenic)	3.0	Undetected	Undetected
F(Fluorine)	3.0	-	0.09
Hg(Mercury)	3.0	Undetected	Undetected
Cr(Chromium)	3.0	Undetected	0.06
Se(Selenium)	3.0	50.8%	0.05

BioSulfa(powder) antibacterial test (Staphylococcus aureus)

한국분석시험연구원
KATR
Korea Analysis Test Researcher

서대문구 연희로 261-28, 2층(홍은동)
T : 1670-9936 F : 02-6016-9711
Homepage : http://katr.re.kr

시험성적서

접수 번호 : KATR180702-001
성적서 번호 : KAAA180715-006
신청 회사 : 에코바이오홀딩스 주식회사

접수 일자 : 2018. 07. 02
발급 일자 : 2018. 07. 15
용도 : 품질관리

주소 : 서울특별시 서초구 서운로26길 5(서초동, 토opil에코빌딩)
담당자 : 한우호
제출처 :
시료명 : 바이오황 #

시험항목	Control	시험결과						
		#1	#2	#3	#4	#5	#6	#7
ASTM E 2149-13a								
<i>Staphylococcus aureus</i> 생균수(CFU/mL)	2.8X10 ⁵	ND	ND	ND	2.5X10 ⁵	1.7X10 ⁴	1.0X10 ⁴	1.1X10 ⁴
ATCC 6538 균감소율(%)		99.9	99.9	99.9	10.7	93.9	96.4	96.1
접종균액의 농도 : 2.5 X 10 ⁵ CFU/mL								


※ 시료 종류 : 분말
 ※ 시료량 : 1.0 g
 ※ 시료 반응 시간 : 1시간
 ※ 균감소율(%) = ((control 균수)-(시료 균수))/(control 균수)X100

한국분석시험연구원

비고 1. 이 성적서는 신청인이 제시한 시료 및 시료명으로 시험한 결과로서 전체 제품에 대한 품질을 보증하지 않습니다.
 2. 이 성적서는 KATR의 동의 없이 홍보, 선전, 광고 및 소송용으로 사용될 수 없으며, 용도 이외의 사용을 금합니다.

99.9% antibacterial effect on *Staphylococcus Aureus*

[2018.07.15] * Analysis: KATR (Korea Institute of Analysis and Testing)



Efficacy of on BioSulfa on chicken mite



Chicken mite adults were killed 45.6%(T0 1: water 2),
Mite eggs were killed 95.0%(T0 1: water 5)

[2018.08.29] Biogenoci farmcq Research & Development Center

시험 성적서

접수일자	2018. 08. 29.	접수번호	01-18-008
시료명	T0	시험대상	<i>Dermanyssus gallinae</i>
제조사	에코바이오텔링스	시험구분	<i>D. gallinae</i> 사멸율 실험
의뢰자	"T0" 시료의 <i>D. gallinae</i> 사멸율 확인	소재지	서울특별시 서초구 서운로 26길 5

귀하께서 우리 팜씨큐 연구개발센터에 의뢰한 시료에 대한 시험결과는 다음과 같습니다.

시험방법 및 결과

시험방법

- 약제노출 후 사멸율 (mortality, %) 측정
- D. gallinae* 성충 100마리, 알 20개 집중, 3회 반복 실험 수행
- 실험농도 : 1:2, 1:5, 1:10
- 약제 농도별 *D. gallinae* 사멸율 (mean±SE, %) 측정

① T0 성충 농도별 사멸율(Mortality, %) 측정 결과

성충 결과 (In vitro)	처리 후 결과 (Mortality ± SE)
T0	24시간 후
1:2	45.6 ± 5.4
1:5	0.8 ± 0.4
1:10	3.1 ± 0.8
Control	0.0 ± 0.0

② T0 알 농도별 사멸율(Mortality, %) 측정 결과

알 결과 (In vitro)	처리 후 결과 (Mortality ± SE)
T0	72시간 후
1:2	85.0 ± 2.9
1:5	95.0 ± 2.9
1:10	45.0 ± 7.6
Control	0.0 ± 0.0

③ 실험 결과 분석


- 의뢰받은 T0 시료에 대한 성충 실험결과, 24시간 후에 1:2 희석배율에서 45.6%의 사멸율을 보였고, 1:10에서는 3.1%의 사멸율을 보였다. 1:2 희석배율을 살펴보면 물보다는 약간 점도가 있었으며, 뿌린 뒤 말랐을 때 간혀죽은 개체가 많았다. 실험자의 의견으로 1:2의 경우 화학적 기작이 아닌 물리적 기작으로 *D. gallinae*를 죽였을 가능성이 높다.
- 알 실험 결과, 1:2배 희석배율에서 85.0%의 사멸율을 보였고, 1:10에서는 45.0%의 사멸율을 보였다. 1:2보다는 1:5에서 더 효과가 좋았다.

비고 : 상기 시험성적서는 의뢰자가 제공한 시료에 대한 결과이며, 시료명은 의뢰자가 제시한 것입니다. 본 성적서는 시험의뢰 목적 이외의 광고, 선전 등 상업적인 용도나 법적인 해결의 용도로 사용될 수 없고 임의로 시험성적서를 변형할 수 없습니다.


2018년 09월 04일

주식회사 바이오지노키 팜씨큐 연구개발센터

Results of mortality for adult mite



Results of mortality for mite egg



Adult result	Result after treatment (Mortality ± SE) (After 24 hours)
1:2	45,6 ± 5,4
1:5	0,8 ± 0,4
1:10	3,1 ± 0,8
Control	0,0 ± 0,0

Egg result	Result after treatment (Mortality ± SE) (After 72 hours)
1:2	85,0 ± 2,9
1:5	95,0 ± 2,9
1:10	45,0 ± 7,6
Control	0,0 ± 0,0

Harmful heavy metals analysis

BEYOND ASIAN HUB, TOWARD GLOBAL WORLD

TEST REPORT

우 22829 인천광역시 서구 가재울로 68 (가좌동) TEL (032)5709-700 FAX (032)575-5613

성적서번호 : TAS-020450 접수 일자 : 2016년 09월 23일
 대표자 : 송효순 시험완료일자 : 2016년 10월 10일
 업체명 : 에코바이오홀딩스(주)
 주소 : 서울특별시 서초구 서운로26길 5(서초동, 토탈에코빌딩)

시료명 : 에코바이오 황

시험결과

시험항목	단위	시료구분	결과치	시험방법
Na	mg/kg	-	22 500	EPA 3050B, 6010D
Pb	mg/L	-	불검출	폐기물공정시험기준 : 2015
Cu	mg/L	-	0.014	폐기물공정시험기준 : 2015
As	mg/L	-	불검출	폐기물공정시험기준 : 2015
Hg	mg/L	-	불검출	폐기물공정시험기준 : 2015
CN ⁻	mg/L	-	불검출	폐기물공정시험기준 : 2015
Cr(VI)	mg/L	-	불검출	폐기물공정시험기준 : 2015
Cd	mg/L	-	불검출	폐기물공정시험기준 : 2015

* 용도 : 품질관리용

비고 : 1. 이 성적서는 의뢰자가 제시한 시료 및 시료명으로 시험한 결과로써 전체 제품에 대한 품질을 보증하지 않으며, 성적서의 진위확인은 홈페이지(www.ktr.or.kr) 또는 QR code로 확인 가능합니다.
 2. 이 성적서는 홍보, 선전, 광고 및 소송용 등으로 사용될 수 없으며, 용도 이외의 사용을 금합니다.
 3. 이 성적서는 원본(동본 포함)만 유효하며, 사본 및 전자 인쇄본/파일본은 결과치 참고용입니다.

Long Jinyoung
 작성자 : 홍진영
 E-mail : longhango@ktr.or.kr

Kim Sun-il
 기술책임자 : 김선일
 Tel : 1577-0091(ARS ①-④)

2016년 10월 10일

KTR 한국화학융합시험연구원

위변조 확인용 QR code

Page : 1 of 1

KTR KOREA TESTING & RESEARCH INSTITUTE KTR-QP-T09-F01-02(07) A4(210 X 297)

Harmful heavy metals not detected

시료명 : 에코바이오 황

시험결과

시험항목	단위	시료구분	결과치	시험방법
Na	mg/kg	-	22 500	EPA 3050B, 6010D
Pb	mg/L	-	불검출	폐기물공정시험기준 : 2015
Cu	mg/L	-	0.014	폐기물공정시험기준 : 2015
As	mg/L	-	불검출	폐기물공정시험기준 : 2015
Hg	mg/L	-	불검출	폐기물공정시험기준 : 2015
CN ⁻	mg/L	-	불검출	폐기물공정시험기준 : 2015
Cr(VI)	mg/L	-	불검출	폐기물공정시험기준 : 2015
Cd	mg/L	-	불검출	폐기물공정시험기준 : 2015

* 용도 : 품질관리용

[2016.10.10] * Analysis : KTR (Korea Testing & Research Institute)

BEYOND ASIAN HUB, TOWARD GLOBAL WORLD

TEST REPORT

우 22829 인천광역시 서구 가재울로 68 (가좌동) TEL (032)5709-700 FAX (032)575-5613

성적서번호 : TAS-020449 접수 일자 : 2016년 09월 23일
 대표 자 : 송효순 시험완료일자 : 2016년 10월 10일
 업체 명 : 에코바이오홀딩스(주)
 주 소 : 서울특별시 서초구 서운로26길 5(서초동, 토탈에코빌딩)
 시 료 명 : 에코바이오 황

시험결과

시험항목	단위	시료구분	결과치	시험방법
S	%	-	50.8	KS M 8088 : 2015(준용)
수분	%	-	49.1	KS M 0010 : 2011

* 용도 : 품질관리용

비 고 : 1. 이 성적서는 의뢰자가 제시한 시료 및 시료명으로 시험한 결과로써 전체 제품에 대한 품질을 보증하지 않으며, 성적서의 진위확인인 홈페이지(www.ktr.or.kr) 또는 QR code로 확인 가능합니다.
 2. 이 성적서는 홍보, 선전, 광고 및 소송용 등으로 사용될 수 없으며, 용도 이외의 사용을 금합니다.
 3. 이 성적서는 원본(등본 포함)만 유효하며, 사본 및 전자 인쇄본/파일본은 결과치 참고용입니다.

Hong Jinyoung Kim Sun-il
 작성자 : 홍진영 기술책임자 : 김선일
 E-mail: longhango@ktr.or.kr Tel : 1577-0091(ARS ①-④)

2016년 10월 10일

KTR 한국화학융합시험연구원 위변조 확인용 QR code

Page : 1 of 1

KTR KOREA TESTING & RESEARCH INSTITUTE KTR-QP-T09-F01-02(07) A4(210 X 297)

well-controlled sulfur content

시 료 명 : 에코바이오 황

시험결과

시험항목	단위	시료구분	결과치	시험방법
S	%	-	50.8	KS M 8088 : 2015(준용)
수분	%	-	49.1	KS M 0010 : 2011

* 용도 : 품질관리용

[2016.10.10] * Analysis: KTR (Korea Testing & Research Institute)

Component Analysis

Crude Proteins (0.56 %), Oleic acid (0.04g/100g) detected

No. : D2018051202

Certificate of Analysis

Date of Application : 2018-05-15	Date of Manufacture :
No. of Sample : D2018051202	Expiration Date :
Lot No. :	
Inspection Purpose : Reference only	
Commodity : 180411_BioSulfa WP	
Applicant	Name : EcoBio Holdings Co., Ltd. Hyo-Soon Song Company address : 06609 TotalEco B/D, 5 Seom-ro 26-gil, Seocho-gu Seoul, South Korea

Analytical Result

Free amino acid(Threonine)(mg/100g)	Not detected
Free amino acid(Cystine)(mg/100g)	Not detected
Free amino acid(Tyrosine)(mg/100g)	Not detected
Free amino acid(Arginine)(mg/100g)	Not detected
Free amino acid(Alanine)(mg/100g)	Not detected
Free amino acid(Proline)(mg/100g)	Not detected
Free amino acid(Lysine)(mg/100g)	Not detected
Free amino acid(Histidine)(mg/100g)	Not detected
Free amino acid(Isoleucine)(mg/100g)	Not detected
Free amino acid(Leucine)(mg/100g)	Not detected
Free amino acid(Methionine)(mg/100g)	Not detected
Free amino acid(Phenylalanine)(mg/100g)	Not detected
Free amino acid(Tryptophan)(mg/100g)	Not detected
Free amino acid(Valine)(mg/100g)	Not detected
Free amino acid(Glutamic Acid)(mg/100g)	Not detected
Free amino acid(Aspartic Acid)(mg/100g)	Not detected
Free amino acid(Serine)(mg/100g)	Not detected
Free amino acid(Glycine)(mg/100g)	Not detected
Crude protein(%)	0.56%
Fructose(mg/g)	Not detected
Lactose(mg/g)	Not detected
Glucose(mg/g)	Not detected
Maltose(mg/g)	Not detected
Sucrose(mg/g)	Not detected

Butyric acid(g/100g)	Not detected
Caproic acid(g/100g)	Not detected
Caprylic acid(g/100g)	Not detected
Capric acid(g/100g)	Not detected
Lauric acid(g/100g)	Not detected
Tridecanoic acid(g/100g)	Not detected
Myristic acid(g/100g)	Not detected
Behenic acid(g/100g)	Not detected
r-Linolenic acid(g/100g)	Not detected
Lignoceric acid(g/100g)	Not detected
Linoleic acid(g/100g)	Not detected
Stearic acid(g/100g)	0.01g/100g
arachidonic acid(g/100g)	Not detected
Arachidic acid(g/100g)	Not detected
Alpha Linolenic Acid(g/100g)	Not detected
Oleic acid(g/100g)	0.04g/100g
Palmitic acid(g/100g)	0.01g/100g
Pentadecanoic acid(g/100g)	Not detected
Erucic acid(g/100g)	Not detected
Myristoleic acid(g/100g)	Not detected
cis-10-Pentadecenoic acid(g/100g)	Not detected
Palmitoleic acid(g/100g)	Not detected
Heptadecanoic acid(g/100g)	Not detected
cis-10-Heptadecenoic acid(g/100g)	Not detected
Elaidic acid(g/100g)	Not detected
Linolelaidic acid(g/100g)	Not detected
cis-11-Eicosenoic acid(g/100g)	Not detected
cis-11,14-Eicosadienoic acid(g/100g)	Not detected
cis-8,11,14-Eicosatrienoic acid(g/100g)	Not detected
cis-11,14,17-Eicosatrienoic acid(g/100g)	Not detected
cis-3,8,11,14,17-Eicosapentaenoic acid(g/100g)	Not detected
heneicosanoic acid(g/100g)	Not detected
cis-13,16-Docosadienoic acid(g/100g)	Not detected
cis-4,7,10,13,16,19-Docosahexaenoic acid(g/100g)	Not detected
Tricosanoic acid(g/100g)	Not detected

Nervonic acid(g/100g)	Not detected
-----------------------	--------------

5 . 29 . 2018

We hereby certify that the above are correct.

Korea Health Supplements Association Sub. Korea Health Supplements Institute

Director : Yang, Joo-Hong *Dr. J. H. Yang*

B-101, Korea Bio Park., 700, Daerangpangyo-ro, Bundang-gu, Seongnam-si, Gyeonggi-do, Republic of Korea

Component Analysis / [2018.05.15] * Analysis : KHSI (Korea Health Supplement Institute)

Pesticide residue and Pathogenic Microorganisms analysis

발급번호 : 18-PPA-7-00098

분석 성적서

① 의뢰인	성명	(주)에코바이오홀딩스	사업자등록번호	229-81-28817
	주소	06809 서울특별시 서초구 서운로28길 5 (서초동) 토탈에코빌딩		
② 의뢰내용	대상 물품명	바이오황 25		
	시험 개요	7항목: 황, 다성분농약, 병원성미생물5종		
	용도	유기농업자재 목록공시(신규신청)		

③ 분석(시험) 성적 :

항목	성적(단위)	비고
주성분(황)	28.20 %	
다성분농약(322성분)	불검출 mg/kg	
<i>E. coli</i> 0157:H7	불검출	
<i>Salmonella</i> spp.(정성)	불검출	
<i>Staphylococcus aureus</i> (정성)	불검출	
<i>Bacillus cereus</i> (정성)	불검출	
<i>Listeria monocytogenes</i> (정성)	불검출	
	이하 여백	

「농업기술실용화재단 분석검정 의뢰 및 처리규정」 제4조의 규정에 의하여 2018년 05월 10일자로 의뢰한 시료에 대한 분석(시험) 성적입니다.

2018년 06월 04일

이 성적은 신청인이 제출한 시료를 분석한 것으로
관련사항 이외의 선전 소송 등 증거자료로 사용하실
수 없습니다.

농업기술실용화재단 이사장

1 / 1 18-PPA-7-00098

Pesticide residue and pathogenic microorganisms not detected

시험 개요	7항목: 황, 다성분농약, 병원성미생물5종	
용도	유기농업자재 목록공시(신규신청)	

분석(시험) 성적 :

항목	성적(단위)	비고
주성분(황)	28.20 %	
다성분농약(322성분)	불검출 mg/kg	
<i>E. coli</i> 0157:H7	불검출	
<i>Salmonella</i> spp.(정성)	불검출	
<i>Staphylococcus aureus</i> (정성)	불검출	
<i>Bacillus cereus</i> (정성)	불검출	
<i>Listeria monocytogenes</i> (정성)	불검출	
	이하 여백	

[2018.06.04] * Analysis : Foundation of Agri. Tech Commercialization AND Transfer


BioSulfa25 efficacy test of Powdery mildew control on Oriental melon (*Cucumis melo L. var. makua*)

T.N: GET1806-15

Trial of organic agricultural materials against Powdery mildew on Oriental melon

KBSI (Korea Bio Safety Institute)

[2018.07.23] Chungbuk Eumseong-gun, Saenggeuk-myeon
82.7% control effect of powdery mildew on Korean melons observed



1. Objectives
 This test is intended to serve as an organic agricultural materials quality certification registration materials for Powdery mildew on Cucumber.

2. Materials and Methods

a. Methods of the phytotoxicity and efficacy testing shall be follow to the notification of RDA(Rural Development Administration)

b. Target diseases : Powdery mildew (*Sphaerotheca fusca*)

c. Host(race) : Oriental melon(chammaseuncheonchamoe)

d. Incidence of target diseases : 20.2% disease severity of average in untreated control was sufficient to review the efficacy.

e. Treatment of chemicals

Chemicals	Content (%)	Efficacy		Phytotoxicity		Sponsor
		Dilution/Dosage	Method	Standard	Double dosage	
Ecobiosulfur25	Sulfur 55.6	1,000	When the initial of diseases occur, to spray 3 times at interval of 10 days (6/12, 6/22, 7/2)	1,000 (6/12)	500 (6/12)	Ecobio holdings
Untreated control	-	-	-	-	-	-

f. Cultural practices : Cultivation under structure. Semiforcing culture. 200×45 interval transplant at 15th February 2018, vinyl mulching. During the test, it was used 'Ecobiosulfur25' chemical except another fungicides.

3. Method of assessment

Division	investigating items	Time of investigation	Date of investigation	Method of investigation
Efficacy	Disease severity	1	7/12	Investigate the disease leaf area on over 100 leaves at 10days after final treatment
Phytotoxicity	Seemingly phytotoxicity existence	3	6/15, 6/17, 6/19	Observe the phytotoxicity at 3, 5, 7days after treatment

4. Results

a. Efficacy

- Trial of efficacy against powdery mildew on Oriental melon (10days after final treatment)

Chemicals	Disease severity(%)				Significant difference (DMRT)	Control value (%)
	I	II	III	Average		
Ecobiosulfur25	4.3	2.8	3.5	3.5	b	82.7
Untreated control	18.8	21.3	20.5	20.2	a	-

C.V.(%) ----- (12.0)

5. Summary

a. Efficacy

- Ecobiosulfur25 exhibited a superior control effect over 82.7% compared to untreated control.

b. Phytotoxicity

- Ecobiosulfur25 were not phytotoxicity in standard and double dosage.

6. Discussion

- The result of this study indicates the Ecobiosulfur25 is available at an organic farming material since there was no phytotoxicity and exhibited a superior control effect over 82.7% compared to untreated control in Oriental melon. So, this is considered to be practical against powdery mildew.


BioSulfa25 efficacy test of red spider mite control on apples

T.N: GET1806-18

Trial of organic agricultural materials against red spider mite in apple orchard

KBSI (Korea Bio Safety Institute)

[2018.07.23] Gyeongsangbuk-do Yecheon-gun Yecheon-eup
52.1% control rate of red spider mites on apples observed



1. Objectives

This is intended to serve as an organic agricultural materials quality certification registration materials for red spider mite in apple orchard.

2. Materials and Methods

a. Methods of the phytotoxicity and efficacy testing shall be follow the procedure given in RDA(Rural Development Administration)

b. Target insects: Red spider mite (*Panonychus ulmi*)

c. Host(race): Apple(Fuji)

e. Incidence of target insects: The 140.0 of target insects in untreated control at before treatment was sufficient to review the efficacy.

f. Treatment of chemicals

Chemicals	Content (%)	Efficacy		Sponsor
		Dilution/Dosage	Method	
Biosulfur25	Sulfur 55.6	1,000	When initial occurrence, to spray 1 time (6/20)	Ecobio Co, Ltd.
Untreated control	-	-	-	-

g. Cultural practices: Cultivation of bare ground. Slender spindle bush form(type). eleven-years. 4×1.5m interval trees. During the test, it was used only 'Biosulfur 25' chemical that except another pesticides.

h. Plot design: Completely randomized design with 3 replication

Division	Treats	Replicates	Total area	Number of tree	Number of trees to use	The total number of trees to use
Efficacy	2	3	6	1	6	6

3. Method of assessment

Division	Investigating items	Time of investigation	Date of investigation	Method of investigation
Efficacy	Survival rate	3	6/12, 6/19, 6/26	Investigate the mites on 30 leaves before treatment and at 7, 14days after treatment

4. Results

a. Efficacy

○ Trial of efficacy against red spider mite in apple orchard(7days after treatment)

Chemicals	Density of before treatment	Survival rate(%)				Significant difference (DMRT)	Control value (%)
		I	II	III	Average		
Biosulfur25	130.0	67.8	44.4	51.1	54.4	b	52.1
Untreated control	140.0	112.5	106.0	122.4	113.6	a	-

C.V.(%) ----- (12.3)

○ Trial of efficacy against red spider mite in apple orchard(14days after treatment)

Chemicals	Density of before treatment	Survival rate(%)				Significant difference (DMRT)	Control value (%)
		I	II	III	Average		
Biosulfur25	130.0	64.4	43.0	48.2	51.9	b	57.8
Untreated control	140.0	119.5	117.4	132.2	123.0	a	-

C.V.(%) ----- (11.1)

5. Summary

○ Biosulfur25 exhibited a superior control effect over 52.1% compared to untreated control.

6. Discussion

○ The result of this study indicates the Biosulfur25 is available at an organic farming material. So, this is considered to be practical against red spider mite.



BioSulfa25 Acute contact toxicity test for honeybees

Study No. : ETBC-18016

Final Report


최종보고서

바이오향25의 꿀벌 (*Apis mellifera*)에 대한
급성접촉독성시험

ETBC-18016

KBSI (Korea Bio Safety Institute)

[2018.06.12] Contact toxicity on honeybee is not detected



7. Tables

Table 1. Cumulative mortality of honeybees

Nominal dose ^a (µg/bee)	No. of exposed honeybees	Cumulative mortality			Mortality (Death / Total)	
		4 hr	24 hr	48 hr	24 hr	48 hr
Untreated control	10	0	0	0	0% (0 / 30)	0% (0 / 30)
	10	0	0	0		
	10	0	0	0		
Negative control ^b	10	0	0	0	0% (0 / 30)	0% (0 / 30)
	10	0	0	0		
	10	0	0	0		
100.000	10	0	0	0	0% (0 / 30)	0% (0 / 30)
	10	0	0	0		
	10	0	0	0		

a: Based on main ingredient input ratio
b: Distilled water+Acetone (8:2), 1 µL/bee

Study No. : ETBC-18016

Final Report

Table 2. Behavioral abnormalities of honeybees

Nominal dose ^a (µg/bee)	No. of exposed honeybees	Abnormal response		
		4 hr	24 hr	48 hr
Untreated control	10	N(10 ^b)	N(10)	N(10)
	10	N(10)	N(10)	N(10)
	10	N(10)	N(10)	N(10)
Negative control ^c	10	N(10)	N(10)	N(10)
	10	N(10)	N(10)	N(10)
	10	N(10)	N(10)	N(10)
100.000	10	N(10)	N(10)	N(10)
	10	N(10)	N(10)	N(10)
	10	N(10)	N(10)	N(10)



a: Based on main ingredient input ratio
b: Number of honeybees
c: Distilled water+Acetone (8:2), 1 µL/bee

※ Observation key

N: Normal
A: Hyperactivity
B: Mobile but not working or flying normally
C: Alive but unable to walk or fly
NA: Not applicable, not observed because of 100% mortality

KBSI (Korea Bio Safety Institute)

[2018.06.12] Oral toxicity on rat is not detected



Final Report

Study No. : ETO-18022

7. Tables [Group summary]

Table 1. Mortality and clinical signs

Group	Dose (mg/kg bw)	Sex	Number of animals	Clinical signs	Mortality	LD ₅₀
1	2000	Female	3	No abnormality detected	0/3 ^a	>2000 ~ ≤5000 mg/kg bw
2	2000	Female	3	No abnormality detected	0/3	

a : Number of dead animals/Number of tested animals

Table 2. Mean body weights

Group	Dose (mg/kg bw)	Sex	Number of animals	Days after administration (g)		
				0	7	14
1	2000	Female	3	184.8 ± 3.4 ^a	219.7 ± 2.1	237.3 ± 8.7
2	2000	Female	3	208.7 ± 7.2	234.6 ± 4.0	254.8 ± 3.6

a : Mean ± standard deviation

KBSI (Korea Bio Safety Institute)


[2018.06.12] Dermal toxicity on rats is not detected

Study No. : ETP-18016 *Final Report*

최종보고서

랫드에 대한 바이오황25의 급성경피독성시험

ETP-18016



Study No. : ETP-18016 *Final Report*

7. Tables [Group summary]

Table 1. Mortality and clinical signs

Group	Dose (mg/kg bw)	Sex	Number of animals	Clinical signs	Mortality (dead / total)	LD ₅₀
1	4000	Male	5	No abnormality detected	0% (0 / 5) ^a	> 4000 mg/kg bw
2	4000	Female	5	No abnormality detected	40% (2 / 5)	

a : Number of Death animals / Number of tested animals



Table 2. Mean body weights

Group	Dose (mg/kg bw)	Sex	Number of animals	Days after administration (g)		
				0	7	14
1	4000	Male	5	260.9 ± 11.5 ^a	304.6 ± 14.4	363.6 ± 13.1
2	4000	Female	5	196.6 ± 9.8	211.0 ± 9.7	230.4 ± 8.8

a : Mean ± standard deviation

KBSI (Korea Bio Safety Institute)

[2018.06.12] Toxicity on freshwater fish is not detected



7. Tables

Table 1. Cumulative mortality of *Oryzias latipes*

Nominal concentration ^a (mg/L)	Number of fish	Cumulative mortality				
		0 hr	24 hr	48 hr	72 hr	96 hr
Control	10	0	0	0	0	0
10.0	10	0	0	0	0	0

a: Based on nominal concentration of main ingredient input ratio

Table 2. Abnormal response of *Oryzias latipes*

Nominal concentration ^a (mg/L)	Number of fish	Abnormal response			
		24 hr	48 hr	72 hr	96 hr
Control	10	NOR(10 ^b)	NOR(10)	NOR(10)	NOR(10)
10.0	10	NOR(10)	NOR(10)	NOR(10)	NOR(10)

a: Based on nominal concentration of main ingredient input ratio

b: Number of fish

※ Observation key

- LOE : Loss of equilibrium
- SUR : Fish mainly at the surface
- HEM : Hemorrhage
- VDE : Vertebral deformation
- BOT : Fish mainly at the bottom
- NOR : Normal
- NA : Not applicable, not observed because of 100% mortality

KBSI (Korea Bio Safety Institute)


[2018.06.12] Muscos membrane irritation on rabbit eye is not detected

Study No. : ETE-18015 *Final Report*

최종보고서

New Zealand White계 토끼에 대한 바이오황25의
안점막자극성시험

ETE-18015



Study No. : ETE-18015 *Final Report*

7. Tables



Table 1. Mortality and clinical signs

Group	No. of treatment	Days after application				Mortality
		0	1	2	3	
No eye washed	1	NOR ^a	NOR	NOR	NOR	0/3 ^b
	2	NOR	NOR	NOR	NOR	
	3	NOR	NOR	NOR	NOR	

a : Normal
b : Number of dead animals/Number of tested animals

KBSI (Korea Bio Safety Institute)

[2018.06.12] Skin irritation on rabbit is not detected



Final Report

Study No. : ETD-18015

Table 3. Evaluation of skin irritation (1/2)

Phases ^a	Number of animals	Sites	Days after treatment			
			0	1	2	3
Erythema & Eschar	1	Control sites	0	0	0	0
		Test sites	0	0	0	0
	2	Control sites	0	0	0	0
		Test sites	0	0	0	0
	3	Control sites	0	0	0	0
		Test sites	0	0	0	0
Edema	1	Control sites	0	0	0	0
		Test sites	0	0	0	0
	2	Control sites	0	0	0	0
		Test sites	0	0	0	0
	3	Control sites	0	0	0	0
		Test sites	0	0	0	0

a : Time after topical treatment

BioSulfa 50% Organic Materials Certification



Listed Number : 1-6-014
 Certification Number : 1-6-002

Quality Certified by the Korean government !

공시번호 : 제 공시-1-6-014호

유기농업자재 공시서

1. 업체명 : 에코바이오홀딩스 (주) Issue number : 18-25

3. 주소(사업장) : 서울특별시 강남구 테헤란로 122 에코비오홀딩스빌딩 1001호

4. 자재의 명칭 : 황

5. 자재의 구분 : 병해충제

6. 상표명 : 에코바이오황

7. 주성분(원료)의 종류 및 - 주성분 : 황 - 원료의 종류 및 함량

8. 유효기간 : 2018.10.11

9. 제조장 주소 : 인천광역시 남동구 대곡로 61

10. 최초 광고일 : 2015.11.11

11. 최초 공시기관 : 농업기

Environment-friendly Agricultural & Organic Inputs Product		National Notice	
Notice Number	National Notice-1-6-014		
Company Name	EcoBio Holdings Co., Ltd.	Chief Executive Officer	Hyo-soon Song
Company Address	5, Seoun-ro 26-gil, Seocho-gu, Seoul, Republic of Korea		
Factory Location	61, Geowol-ro, Seo-gu, Incheon, Republic of Korea		
Organic Inputs Type	An organic agriculture material to control diseases and pests		
Brand name	Eco Bio Sulfur		
Applied Crop	Red pepper, Lettuce, Chinese cabbage, Soybean, Cucumber		
Applied Pest	-		
Expiration Date	Oct. 10. 2018. ~ Oct. 09. 2021.		

「친환경농업 육성 및 제2차 및 「농림축산식품부」 지원에 관한 법률 시행규칙에 따라 공시임을 증명합니다.

Date of Issue : Sep. 12. 2018.

Foundation of Agri. Tech. Commercialization & Transfer

품질인증번호 : 제 품질인증-1-6-002호

유기농업자재 품질인증서

1. 업체명 : 에코바이오홀딩스 (주) Issue number : 18-14

3. 주소(사업장) : 서울특별시 강남구 테헤란로 122 에코비오홀딩스빌딩 1001호

4. 자재의 명칭 : 황

5. 자재의 구분 : 병해충제

6. 상표명 : 에코바이오황

7. 주성분(원료)의 종류 및 - 주성분 : 황 50% - 원료의 종류 및 함량

8. 유효기간 : 2016.06.30

9. 제조장 주소 : 인천광역시 남동구 대곡로 61

10. 최초 광고일 : 2015.11.11

11. 최초 공시기관 : 농업기

Environment-friendly Agricultural & Organic Inputs Product		Quality Certification	
Notice Number	Quality Certification-1-6-002		
Company Name	EcoBio Holdings Co., Ltd.	Chief Executive Officer	Hyo-soon Song
Company Address	5, Seoun-ro 26-gil, Seocho-gu, Seoul, Republic of Korea		
Factory Location	61, Geowol-ro, Seo-gu, Incheon, Republic of Korea		
Organic Inputs Type	An organic agriculture material to control diseases and pests		
Brand name	Eco Bio Sulfur		
Applied Crop	Red pepper, Lettuce, Chinese cabbage, Soybean, Cucumber, Strawberry		
Applied Pest	Cucumber(Powdery mildew), Strawberry(Two spotted spider mite)		
Expiration Date	June. 30. 2016. ~ June. 29. 2019.		

「친환경농업 육성 및 제2차 및 「농림축산식품부」 지원에 관한 법률 시행규칙에 따라 품질인증을 증명합니다.

Date of Issue : April. 10. 2018.

Foundation of Agri. Tech. Commercialization & Transfer


BioSulfa 25% Organic Materials Certification



Listed Number : 1-6-030

Quality Certified by the Korean government !

Issue number : 18-24

Environment-friendly Agricultural & Organic Inputs Product		National Notice	
Notice Number	National Notice-1-6-030		
Company Name	EcoBio Holdings Co., Ltd.	Chief Executive Officer	Hyo-soon Song
Company Address	5, Seoun-ro 26-gil, Seocho-gu, Seoul, Republic of Korea		
Factory Location	61, Geowol-ro, Seo-gu, Incheon, Republic of Korea		
Organic Inputs Type	An organic agriculture material to control diseases and pests		
Brand name	Biosulfur25		
Applied Crop	Red pepper, Lettuce, Grass, Soybean, Cucumber		
Applied Pest	-		
Expiration Date	Aug. 23. 2018. ~ Aug. 22. 2021.		
<p>In accordance with Article 37 of 「ENVIRONMENT-FRIENDLY AGRICULTURE PROMOTION ACT」, I hereby certify that the product above is listed on the National Notice List of Environment-friendly Agricultural & Organic Inputs in the Republic of Korea.</p>			
<p>Date of Issue : Sep. 12. 2018.</p>			
<p>Foundation of Agri. Tech. Commercialization & Transfer</p> 			





BioSulfa

Crop test results

Test results on various crops as well as observations notes.

Benefits of utilizing Sulfur in agriculture

- Healthy crops due to immunity improvement
- Minimization of the reliance on protective agent
- Increase of yield (Increase of organic matter content)
- Universalization of eco-friendly agricultural techniques (Environment improvement)



High Value
“BioSulfa”




Summary of BioSulfa efficacy on Powdery Mildew and Mites




■ No treatment ■ Treatment



Powdery mildew



Mites



¹⁾ Cucumber (2016.04.08~2016.05.29) ²⁾ Oriental melon (2018.06.12~2018.07.12)

¹⁾ Strawberry (2016.04.26~2016.05.10) ²⁾ Apple (2018.06.12~2018.07.04)

Certified Institute Test of BioSulfa on Powdery Mildew and Mites(1/4)

Performed by the Korea Bio-Safety Institute / Cucumber 2016.04.08~05.08 (Ipjang), 2016.04.29~05.29 (Namseon)

Crop	Targets	Results	Treatment Images	
<p>Cucumber</p>	<ul style="list-style-type: none"> Phytotoxicity Powdery Mildew (<i>Sphaerotheaca fusca</i>) 	<ul style="list-style-type: none"> Standard dilution(x1,000) and double dosage dilution(x500) were applied 3 times Foliar Spraying after breakout in every 10 days 85% efficacy on powdery mildew at Ipjang farm 83.6% efficacy on powdery mildew at Namseon farm Greenhouse conditions Test was carried out at 3 different region at same time during 30 days No phytotoxicity in standard dose and double dosage. 		
			<p>< test photo 1 ></p>	<p>< Test photo 2 ></p>
			<p>< test photo 3 ></p>	<p>< Phytotoxicity test result > (3, 5, 7 days after treatment)</p>





Certified Institute Test of BioSulfa on Powdery Mildew and Mites(2/4)

Performed by the Korea Bio-Safety Institute / 2018.06.12-2018.07.12 (Namseon, Saengguk)

Crop	Targets	Results	Treatment Images							
<p>Oriental melon</p>	<ul style="list-style-type: none"> • Phytotoxicity • Powdery Mildew (<i>Sphaerotheaca fusca</i>) 	<ul style="list-style-type: none"> • Standard dilution(x1,000) and double dosage dilution(x500) were applied • 3 times Foliar Spraying after breakout in every 10 days • 78.4% efficacy on powdery mildew at Namseon farm • 82.7% efficacy on powdery mildew at Saengguk farm • Greenhouse conditions • Test was carried out at 3 different region at same time during 30 days • No phytotoxicity in standard dose and double dosage. 			<p>< test photo 1 ></p>	<p>< Test photo 2 ></p>			<p>< test photo 3 ></p>	<p>< Phytotoxicity test result > (3, 5, 7 days after treatment)</p>





Certified Institute Test of BioSulfa on Powdery Mildew and Mites(3/4)

Performed by the Korea Bio-Safety Institute / 2016.04.26~2016.05.10 (Gangok)


Crop	Targets	Results	Treatment Images
<p>Strawberry</p>	<ul style="list-style-type: none"> • Phytotoxicity • Mite (<i>Tetranychus urticae</i>) 	<ul style="list-style-type: none"> • Standard dilution(x1,000) and double dosage dilution(x500) were applied • 1 time Foliar Spraying after breakout • 65.9% efficacy on strawberry mite at Gangok farm • Test was carried out at 3 different region at same time during 14 days • No phytotoxicity in standard dose and double dosage 	<div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">  <p>< test photo 1 ></p> </div> <div style="text-align: center;">  <p>< Test photo 2 ></p> </div> </div> <div style="display: flex; justify-content: space-around; margin-top: 20px;"> <div style="text-align: center;">  <p>< test photo 3 ></p> </div> <div style="text-align: center;">  <p>< Phytotoxicity test result > (3, 5, 7 days after treatment)</p> </div> </div>

Certified Institute Test of BioSulfa on Powdery Mildew and Mites(4/4)

Performed by the Korea Bio-Safety Institute / 2018.06.12-2018.06.26 (Yecheon), 2018.06.20-2018.07.04 (Gangmok)

Crop	Targets	Results	Treatment Images
<p>Apple</p>	<ul style="list-style-type: none"> Phytotoxicity Mite (<i>Panonychus ulmi</i>) 	<ul style="list-style-type: none"> Standard dilution(x1,000) and double dosage dilution(x500) were applied 1 time Foliar Spraying after breakout 57.8% efficacy on apple mite at Yecheon farm 62.1% efficacy on apple mite at Gangmok farm Test was carried out at 3 different region at same time during 14 days No phytotoxicity in standard dose and double dosage. 	<div style="display: flex; flex-direction: column; align-items: center;"> <div style="display: flex; justify-content: space-around; width: 100%;"> <div style="text-align: center;">  <p>< test photo 1 ></p> </div> <div style="text-align: center;">  <p>< Test photo 2 ></p> </div> </div> <div style="display: flex; justify-content: center; margin-top: 20px;"> <div style="text-align: center;">  <p>< test photo 3 ></p> </div> <div style="text-align: center; margin-left: 20px;"> <p style="font-size: small;">사과(홍묘) 약해조사(약제처리 후 3, 5, 7일차)</p>  <p>< Phytotoxicity test result > (3, 5, 7 days after treatment)</p> </div> </div> </div>


Biosulfa test on oligonychus punicae control for avocado



	ADA	3DDA	7DDA	13DDA
BioSulfa	49.38	5.13	2.25	0.65
SULPHA-N	43.13	23.38	17.50	12.25
OMEX SW7	43.13	6.00	2.50	0.25
SILICATO K	50.00	50.00	50.00	50.00

- ◆ Test performed with **BioSulfa**, SULFA-N, OMEX SW7, and SILICATO K
- ◆ **BioSulfa** presented significantly higher efficacy compared to other products

Biosulfa test on Phyllocoptruta oleivora control forcitrus



- ◆ Test performed with **BioSulfa**, SULFA-N, OMEX SW7, and SILICATO K
- ◆ **BioSulfa** presented significantly higher efficacy compared to other products

Biosulfa test for *Phyllocoptruta oleivora* control on citrus

PRODUCTOS	Nº PLANTAS	Nº FRUTOS	Nº ACAROS ADA	Nº ACAROS DDA
			27/04/2018	30/04/2018
BioSulfa	P1	F1	64	6
		F2	65	4
		F3	38	5
	P2	F4	64	8
		F5	28	4
		F6	72	4
OMEX SW7	P1	F1	48	32
		F2	37	15
		F3	26	12
	P2	F4	53	32
		F5	33	32
		F6	46	12

PRODUCTOS	Nº PLANTAS	Nº FRUTOS	Nº ACAROS ADA	Nº ACAROS DDA
			27/04/2018	30/04/2018
SULFA N	P1	F1	56	32
		F2	34	30
TESTIGO	P2	F3	16	00
		F4	25	25
	P1	F1	43	50
		F2	52	55
P2	F3	47	71	
	F4	41	62	

- ◆ Test performed with **BioSulfa**, SULFA-N, OMEX SW7, and SILICATO K
- ◆ **BioSulfa** presented significantly higher efficacy compared to other products

BioSulfa Application Example(Soil disinfection)



Lee Chang Hee (Gwangju, Gyeonggi) 2014. 04. 02
- BioSulfa was added to soil
- No effect was observed on other microorganisms.

Yang Jae Hwa (Mungyeong, Gyeongbuk) May 26, 2016
- x500 dilution was applied for sterilization
- x2,000 dilution was applied after planting



Kim Tae-Hwan (Nyongju, North Gyeongsang Province) 27.6.7
- Overeatic Soil Sterilization



Park Jung-il (Ulsan, Gyeongnam) Nov. 16, 2016
- Sterilization after Typhoon Chava

BioSulfa Application example(Winter control)



Kim Seon-mi (Gimcheon, Gyeongbuk)
- For winter control



Kim Hae-soon (Yeongcheon, Gyeongbuk) 2014. 04. 04
- x200 dilution was applied



Kim Sun-mi (Kimcheon, North Gyeongsang Province)
- Biohwang Winter Protection



Jung Moon-seok (Boeeun, North Chungcheong Province) 2016-09
Using a mixture of 500 litres of water and 1 litre of biohwang.
- 2 litres of caterpillar


BioSulfa Application example(Foliar spray application)



Kim Jong-sam (Gohung, Jeonnam) 2016. 06.27
- x200 dilution was applied
- Mite was exterminated



Son Woo Hyun (Cheongsong, Gyeongbuk)
- x250 dilution was applied
- No other fertilizer and pesticide were used



Kim Hakkwon 2017.7.27
- x1,000 dilution was applied to the second year of Fuji,
- x500 dilution was applied to fruit and the 2nd year old, 10 years old trees
- x500 dilution was applied to the fruit and the 15th years old trees
- X125 dilution was applied to fruit and bud(3 years Hongro)

High Kyung Tae (Cheongsong, Gyeongbuk)
- Black-spot disease and rust-resistant noise

BioSulfa Application Example (Vegetable leaf)



Shin Tae-soo (Gohan, Jeonnam) 2014. 04. 26

- x360 dilution was applied
- 2 times for garlic, 3 times for potato

Choi Sung-ryeol (Daejeon) 2016. 06. 12

- x500 dilution applied for ginseng



Lee Young-seok (Haenam, Jeonnam) 2016. 10 16

- Mixture of BioSulfa + rooting agent + calcium + minor elements was applied 3 times

Kim Young Soo (Gosung, Gyeongnam) 2016. 11. 28

- x2,500 was applied

BioSulfa Application example(Sulfur containing rice)



Park Seok-joon (Hampyeong, Jeonnam) 2017. 07. 23
 - x30 dilution was applied
 - Foliar application

Choi Sun-joon (Gwangyang, Jeonnam) 2016. 08. 10
 - Scatter 500,333,250 times biohwang.
 - Growing eco-friendly organic rice



Lee Chang-min (Yeongam, Jeonnam) 2016. 08. 14
 - BioSulfa treatment

BioSulfa Application Example (Schizandra cultivation)



Gwang-Hyun Lee (Inje, Gangwon) 2016. 09. 14

- BioSulfa was applied 4 times
- No disease was found

Kwon Hyuk-beom (Pyeongchang, Gangwon)



Wi Kwang Hyun (Gangwon Inje) 2016

- Scatter biohwang 4
- No disease detected

Kim, Gyeong-rye (Chungju, Chungbuk) 2017. 09. 07

- Increase yield and harvest without disease

BioSulfa Application example(Walnut tree, banana disaster)



Kang Byung-taek (Youngdong, Chungbuk) May 11, 2015
- Winter sterilization for walnut farm



Kwon-Bae (Jeju) 2017.9.14
- Prevention of banana disease
- BioSulfa was applied on banana every few days

BioSulfa Application example(Apple, Tangerine)



Kim Hak Kwon 2015. 07. 27 / 2016. 08. 20

- 1000 times the amount of biohwang, 500 times the number of beans, 250 times the birth date of 15 years



Go Dong-hee (Jeju) Aug. 20, 2015

- x2,000 dilution was applied to Karahyang
- Spraying every 5 days

Ko Kyung Tae 2016. 08. 09

- Biohwang 3rd spraying
- Grow pesticide free oil

BioSulfa Application example(Powdery mildew)



Jeong Yong-yong (Jinju, Gyeongnam) Dec. 12, 2017
- x2,000 dilution was applied



Photo after treatment



Son Byung-gwan (Iksan, Jeonbuk) 2015. 07. 21
- x2,000 dilution was applied and good effect was found

Photo after 10 days

BioSulfa application case(Anthrax)



Mo Chung Yong (Muan, Jeonnam) 2017. 09. 16
- Red pepper anthrax and strawberry anthrax

Jeong Tae-Mook 2017.7.21
- Green, non-farm pesticide raspberries
- Prevention of biohwang anthrax



Lee Young-Suk (Hannam, Jeonnam)
- 200 times the amount of biohwang in anthrax

BioSulfa Application Example(Mite)



Mo Cheong-yong (Muan, Jeonnam)

- Spotted mites in cucumber
- x10 dilution was applied
- No mite was observed after treatment

Lee Jae-ryong 2016.4.5

- Using biosulfa



Kim Jong-sam 2016.6.27

- Prevention of yuja tree mites
- 200 times biohwang to remove condensation

Jeong Woo-cheol (Gongju, Chungnam) 2017. 06. 19

- Using biosulfa to treat mites

Thank You !!



ECOBIO HOLDINGS CO., LTD.

Total Eco Bldg. 5, Seoun-ro 26-gil, Seocho-gu, Seoul, Republic of Korea
Phone : +82)2-3483-2900 Fax : +82)2-3483-2929
Email : biosulfa@ecobio.co.kr homepage : www.ecobio.co.kr